



INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)

An International Open Access, Peer-reviewed, Refereed Journal

End-Stage Renal Disease In India: A Comprehensive Review Of Etiopathology, Clinical Presentation, And Management Approaches

Dr. Zubair Ali Khan MD (Hom.)

Assistant Professor

Department of Anatomy

Ananya College of Homoeopathy, KIRC, kalol, Gujarat (India)

Abstract

End-stage renal disease (ESRD) is the terminal stage of chronic kidney disease, requiring dialysis or kidney transplantation for survival. In India, rising rates of diabetes and hypertension, along with limited access to care, have made ESRD a growing public health challenge. While modern medicine remains the primary treatment approach through renal replacement therapies and supportive care, alternative systems like homeopathy are gaining interest. Homeopathy, with its individualized, symptom-focused approach, aims to relieve ESRD symptoms such as fatigue, edema, and nausea, and support overall well-being. Emerging evidence and case reports suggest homeopathy may complement conventional care, particularly in improving quality of life and symptom management.

Index terms- ESRD, CKD, homoeopathy, CKI

Introduction

End-stage renal disease (ESRD), also known as stage 5 chronic kidney disease (CKD), represents the final stage of kidney failure, where kidney function declines to below 10-15% of normal. Patients with ESRD require renal replacement therapy (RRT), including dialysis or kidney transplantation, to survive. With a rising incidence globally, ESRD has become a significant public health concern, especially in low- and middle-income countries like India, where early detection and treatment are often delayed.

India is witnessing a rapid increase in ESRD due to the burgeoning rates of diabetes, hypertension, and an aging population. The lack of widespread screening, limited infrastructure, and economic constraints further compound the challenge. This article delves into the etiopathology, risk factors, clinical presentation, diagnostic criteria, and management of ESRD, with a specific focus on the Indian healthcare landscape and the roles of modern and alternative medicine.

Etiopathology

The pathogenesis of ESRD typically involves a progressive decline in glomerular filtration rate (GFR) due to damage to the nephron structure. The etiologies of nephron damage vary but converge on a final common pathway characterized by glomerulosclerosis, tubular atrophy, interstitial fibrosis, and chronic inflammation.

The primary pathological mechanisms include:

1. **Hyperfiltration Injury:** Seen in diabetes and obesity, increased glomerular pressure leads to structural damage.
2. **Immune-mediated Damage:** Conditions like lupus nephritis and glomerulonephritis cause immune complex deposition and inflammation.
3. **Ischemic Injury:** Hypertension causes vascular changes leading to ischemia and scarring.
4. **Toxic Injury:** Long-term use of NSAIDs or exposure to heavy metals can cause direct tubular toxicity.

Predisposing Factors

Several risk factors contribute to the development and progression of ESRD:

1. **Diabetes Mellitus:** The leading cause of ESRD in India, accounting for 30-40% of cases.
2. **Hypertension:** A significant risk factor contributing to nephrosclerosis.
3. **Glomerulonephritis:** Including focal segmental glomerulosclerosis, IgA nephropathy, and membranous nephropathy.
4. **Polycystic Kidney Disease:** A genetic disorder causing progressive cyst formation.
5. **Recurrent Urinary Tract Infections and Obstruction:** Leading to chronic pyelonephritis.
6. **Prolonged Use of Nephrotoxic Drugs:** NSAIDs, aminoglycosides.
7. **Low Birth Weight and Intrauterine Growth Retardation:** Associated with lower nephron numbers.

Prevalence and Current Scenario in India

The Indian CKD registry reports an increasing prevalence of CKD, with an estimated 17% of the adult population showing signs of kidney dysfunction [1]. Of these, approximately 2.2 lakh patients progress to ESRD annually, though fewer than 20% receive RRT [2].

Barriers to treatment include:

- Lack of awareness and routine screening.
- High out-of-pocket expenses.
- Inadequate public sector infrastructure.
- Urban-rural healthcare disparity.

The Indian government has implemented programs such as the Pradhan Mantri National Dialysis Programme (PMNDP) to address this burden, but challenges remain in equitable access and quality of care.

Signs and Symptoms

The clinical manifestations of ESRD are often non-specific and result from the accumulation of toxins, fluid overload, and hormonal dysregulation.

- **General:** Fatigue, weakness, anorexia, weight loss.
- **Cardiovascular:** Hypertension, congestive heart failure, pericarditis.
- **Gastrointestinal:** Nausea, vomiting, metallic taste, gastrointestinal bleeding.
- **Neurological:** Confusion, seizures, peripheral neuropathy.
- **Dermatological:** Pruritus, pallor, uremic frost.

- **Endocrine:** Amenorrhea, infertility, sexual dysfunction.
- **Skeletal:** Bone pain, fractures (renal osteodystrophy).

Complications

ESRD affects nearly every organ system and leads to various complications:

1. **Anemia:** Due to erythropoietin deficiency.
2. **Mineral Bone Disease:** Secondary hyperparathyroidism, osteitis fibrosa.
3. **Cardiovascular Disease:** Leading cause of death in ESRD patients.
4. **Hyperkalemia:** Life-threatening arrhythmias.
5. **Metabolic Acidosis:** Muscle wasting, bone demineralization.
6. **Infections:** Due to immunosuppression.
7. **Malnutrition:** Common and associated with increased mortality.

Diagnostic Criteria

Diagnosis of ESRD is based on clinical evaluation, laboratory tests, and imaging. Key diagnostic criteria include:

1. **Glomerular Filtration Rate (GFR):** $\text{GFR} < 15 \text{ mL/min/1.73 m}^2$ (CKD-EPI or MDRD formula).
2. **Elevated Serum Creatinine and Blood Urea Nitrogen (BUN).**
3. **Electrolyte Imbalance:** Hyperphosphatemia, hyperkalemia, metabolic acidosis.
4. **Imaging:** Ultrasound showing small, echogenic kidneys.
5. **Urinalysis:** Proteinuria, hematuria, cellular casts.
6. **Renal Biopsy:** In selected cases to identify specific etiology.

PARAMETER	DIAGNOSTIC VALUE
GFR	$< 15 \text{ mL/min/1.73 m}^2$
Serum Creatinine	$> 5.0\text{--}6.0 \text{ mg/dL}$ (often higher)
BUN	$> 100 \text{ mg/dL}$
Urine Output	Oliguria ($< 400 \text{ mL/day}$) or Anuria ($< 100 \text{ mL/day}$)
Potassium	$> 5.5 \text{ mEq/L}$ (risk of cardiac arrest if > 6.5)
Phosphate	$> 5.5 \text{ mg/dL}$
Calcium	$< 8.5 \text{ mg/dL}$ (corrected)
Bicarbonate	$< 22 \text{ mmol/L}$ (metabolic acidosis)
Hemoglobin	$< 10 \text{ g/dL}$ (often $< 8 \text{ g/dL}$ in ESRD)
Ultrasound Kidney Size	$< 9 \text{ cm}$; echogenic with cortical thinning

PARAMETER	DIAGNOSTIC VALUE
Urinalysis	Proteinuria, broad/waxy casts, low spec. gravity

Modern Medical Approach

1. Renal Replacement Therapy (RRT)

- **Hemodialysis (HD):** Most common modality. Requires vascular access (AV fistula). Thrice-weekly sessions are standard.
- **Peritoneal Dialysis (PD):** Home-based option, less commonly used in India.
- **Kidney Transplantation:** Gold standard for eligible patients. Challenges include donor shortage and financial constraints.

2. Pharmacological Management

- **Erythropoiesis-stimulating agents:** For anemia.
- **Phosphate binders and Vitamin D analogs:** To manage mineral bone disorder.
- **Diuretics:** For volume overload.
- **Antihypertensives:** ACE inhibitors/ARBs.
- **Sodium bicarbonate:** To correct acidosis.

3. Nutritional Support

- Low-protein diet (0.6–0.8 g/kg/day).
- Sodium and potassium restriction.
- Adequate caloric intake to prevent muscle wasting.

4. Government Initiatives

- **PMNDP:** Free dialysis in district hospitals.
- **Ayushman Bharat – PM-JAY:** Covers up to Rs 5 lakh per family annually for ESRD treatment.

Alternative Medicine Approach

1. Ayurveda

- **Herbal Remedies:** *Punarnava (Boerhavia diffusa)*, *Gokshura (Tribulus terrestris)*.
- **Panchakarma Therapies:** Detoxification and rejuvenation.
- **Rasayana Therapy:** Enhances body resistance.
- Studies suggest potential in delaying progression in early CKD, but limited data in ESRD [3].

2. Yoga and Naturopathy

- **Asanas and Pranayama:** Improve blood pressure control, reduce stress.
- **Detoxifying Diets:** Plant-based, anti-inflammatory.
- Studies show improved quality of life in dialysis patients practicing yoga [4].

3. Integrative Medicine

- AIIMS and CCRAS initiatives exploring integrative models.
- Evidence-based integration may improve symptom control and patient satisfaction.

4. Homeopathic Approach

The goal in ESRD is to:

- Slow down disease progression (in early CKD stages).
- Relieve specific symptoms (e.g., fatigue, edema, nausea, itching).
- Support the patient's overall well-being alongside conventional treatments.

In homeopathy, treatment is Individualized. Practitioners take into account the patient's constitution, mental and emotional state, as well as physical symptoms.

Some efficient work has been done in cases of End stage renal diseases in homeopathy.

Some Indicated homoeopathic remedies[16] in ESRD are as follows-

1- ***Serum Anguillae (Eel Serum)***- Beneficial in Hypertension with oedema of kidney origin. Indicative in Albuminuria, oliguria, dyspnoea, threatening uraemia, hence proves a clinical Renal remedy.

2- ***Terebinthinae Oleum (Turpentine)***- Albuminuria in end stages and few casts. Anasarca in organic kidney diseases. Urine is cloudy and smoky with odour of violets.

3- ***Urea Pura (Carbamide)***- General intoxication from renal disease accompanying diabetes, uraemia, albuminuria and dropsy.

4- ***Juniperus Virginianus (Red Cedar)***- Dropsy of the aged people with suppression of urination. Chronic pyelitis.

5- ***Quebracho (Aspidosperma)***- Uraemic Dyspnoea.

6- ***Apocynum cannabinum (Indian Hemp)***- Diminished excretions. Dropsy with thirst from organic diseases after typhoid, typhus, or abuse of quinine.

7- ***Morphinum*** – Acute as well as chronic uraemia. Cheyne stokes respiration. Tachycardia alternates with bradycardia.

Conclusion

ESRD is a growing public health crisis in India, demanding a multi-faceted approach. While modern medicine provides life-sustaining options like dialysis and transplantation, the integration of evidence-based alternative therapies can enhance patient outcomes and quality of life. Public awareness, early detection, policy support, and inclusive healthcare systems are vital to curbing the ESRD epidemic in India.

References

1. Rajapurkar, M.M., et al. (2012). What do we know about chronic kidney disease in India: First report of the Indian CKD registry. *BMC Nephrology*, 13(10). <https://doi.org/10.1186/1471-2369-13-10>
2. Modi, G.K., & Jha, V. (2006). The incidence of end-stage renal disease in India: A population-based study. *Kidney International*, 70(12), 2131–2133.
3. Gokarn, R.A., et al. (2011). Evaluation of an Ayurvedic formulation in chronic kidney disease: A pilot study. *AYU*, 32(3), 314–320.
4. Satish, M., et al. (2021). Effect of integrated yoga therapy on quality of life in hemodialysis patients: A randomized control trial. *Complement Ther Med*, 58, 102687.
5. Ministry of Health and Family Welfare. (2023). Pradhan Mantri National Dialysis Programme Guidelines. <https://nhm.gov.in>
6. NOTTO (2024). Annual Report 2023-24. <https://notto.mohfw.gov.in/>
7. **KDIGO (2012)**. *Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease*. *Kidney Int Suppl*. 2013;3(1):1-150.
8. **National Kidney Foundation (NKF)**. *KDOQI Clinical Practice Guidelines for Chronic Kidney Disease*. *Am J Kidney Dis*. 2002;39(2 Suppl 1):S1-266.
9. Levey AS, et al. (2003). A more accurate method to estimate GFR from serum creatinine. *Ann Intern Med*. 139(2):137–147.
10. Taal MW, Chertow GM, Marsden PA, et al. (Eds.). (2019). *Brenner & Rector's The Kidney* (11th ed.). Elsevier.
11. Jameson JL, Fauci AS, Kasper DL, et al. (Eds.). (2018). *Harrison's Principles of Internal Medicine* (20th ed.). McGraw-Hill.
12. **KDIGO (2017)**. *CKD-MBD Guideline Update*. *Kidney Int Suppl*. 7(1):1–59.
13. **KDIGO (2012)**. *Anemia in CKD Guidelines*. *Kidney Int Suppl*. 2(4):279–335.
14. Hricak H, Cruz C, Romanski R, et al. (1982). Renal parenchymal disease: sonographic-histologic correlation. *Radiology*. 144(1):141–147.
15. Fogazzi GB, Garigali G. (2003). Urinalysis: Core curriculum 2003. *Am J Kidney Dis*. 42(4):823–833.
16. Boericke W. Pocket Manual of Homoeopathic Materia Medica and Repertory. B Jain Publishers. 9th Edition, 2012, India.